

3. 部位別登録数と割合

3-1. 部位別登録数と割合

| 診断年 | 2010年 | | 2011年 | | 2012年 | | 2013年 | | 2014年 | |
|--------|-------|---------|-------|---------|-------|---------|-------|---------|-------|---------|
| | 件数 | (%) | 件数 | (%) | 件数 | (%) | 件数 | (%) | 件数 | (%) |
| 口腔・咽頭 | 19 | (1.8%) | 27 | (2.3%) | 26 | (2.2%) | 20 | (1.7%) | 27 | (2.4%) |
| 食道 | 21 | (2.0%) | 15 | (1.3%) | 23 | (2.0%) | 23 | (1.9%) | 24 | (2.2%) |
| 胃 | 178 | (16.6%) | 188 | (15.9%) | 180 | (15.4%) | 204 | (17.3%) | 153 | (13.8%) |
| 結腸 | 102 | (9.5%) | 139 | (11.7%) | 140 | (12.0%) | 134 | (11.4%) | 110 | (9.9%) |
| 直腸 | 72 | (6.7%) | 83 | (7.0%) | 60 | (5.1%) | 63 | (5.3%) | 80 | (7.2%) |
| 肝臓 | 51 | (4.8%) | 51 | (4.3%) | 50 | (4.3%) | 49 | (4.2%) | 49 | (4.4%) |
| 胆嚢・胆管 | 31 | (2.9%) | 25 | (2.1%) | 28 | (2.4%) | 23 | (1.9%) | 28 | (2.5%) |
| 膵臓 | 34 | (3.2%) | 38 | (3.2%) | 44 | (3.8%) | 43 | (3.6%) | 47 | (4.2%) |
| 喉頭 | 6 | (0.6%) | 7 | (0.6%) | — | — | 10 | (0.8%) | 6 | (0.5%) |
| 肺 | 55 | (5.1%) | 78 | (6.6%) | 80 | (6.8%) | 76 | (6.4%) | 56 | (5.0%) |
| 骨・軟部 | — | — | 0 | (0.0%) | — | — | — | — | — | — |
| 皮膚 | 43 | (4.0%) | 33 | (2.8%) | 43 | (3.7%) | 43 | (3.6%) | 58 | (5.2%) |
| 乳房 | 59 | (5.5%) | 84 | (7.1%) | 97 | (8.3%) | 84 | (7.1%) | 77 | (6.9%) |
| 子宮頸部 | 28 | (2.6%) | 44 | (3.7%) | 50 | (4.3%) | 44 | (3.7%) | 34 | (3.1%) |
| 子宮体部 | 15 | (1.4%) | 10 | (0.8%) | 19 | (1.6%) | 18 | (1.5%) | 23 | (2.1%) |
| 子宮 | 0 | (0.0%) | 0 | (0.0%) | 0 | (0.0%) | 0 | (0.0%) | 0 | (0.0%) |
| 卵巣 | 11 | (1.0%) | 13 | (1.1%) | 6 | (0.5%) | — | — | 5 | (0.5%) |
| 前立腺 | 138 | (12.9%) | 137 | (11.6%) | 108 | (9.2%) | 103 | (8.7%) | 94 | (8.5%) |
| 膀胱 | 39 | (3.6%) | 39 | (3.3%) | 43 | (3.7%) | 42 | (3.6%) | 54 | (4.9%) |
| 腎・尿路 | 30 | (2.8%) | 25 | (2.1%) | 20 | (1.7%) | 40 | (3.4%) | 32 | (2.9%) |
| 脳・神経 | 41 | (3.8%) | 35 | (3.0%) | 30 | (2.6%) | 34 | (2.9%) | 34 | (3.1%) |
| 甲状腺 | 12 | (1.1%) | 11 | (0.9%) | 11 | (0.9%) | 10 | (0.8%) | 12 | (1.1%) |
| 悪性リンパ腫 | 32 | (3.0%) | 37 | (3.1%) | 37 | (3.2%) | 31 | (2.6%) | 32 | (2.9%) |
| 多発性骨髄腫 | 9 | (0.8%) | 9 | (0.8%) | 10 | (0.9%) | 7 | (0.6%) | 12 | (1.1%) |
| 白血病 | 19 | (1.8%) | 18 | (1.5%) | 13 | (1.1%) | 22 | (1.9%) | 20 | (1.8%) |
| 他の血液 | 10 | (0.9%) | 12 | (1.0%) | 12 | (1.0%) | 15 | (1.3%) | 18 | (1.6%) |
| その他 | 17 | (1.6%) | 26 | (2.2%) | 31 | (2.7%) | 37 | (3.1%) | 25 | (2.3%) |
| 合計 | 1,073 | | 1,184 | | 1,168 | | 1,180 | | 1,111 | |

参考 結腸・直腸の合計を『大腸』、子宮頸部・子宮体部・子宮の合計を『子宮』、悪性リンパ腫・多発性骨髄腫・白血病・他の血液腫瘍の合計を『血液腫瘍』とした場合の上位5部位

